Ester-C® Complex

Vitamin C as Calcium Ascorbate with Citrus Bioflavonoids

DESCRIPTION

Ester-C® Complex capsules, provided by Douglas Laboratories®, contain a non-acidic, calcium ascorbate form of vitamin C combined with a citrus bioflavonoid complex.

Ester-C® is a unique form of vitamin C that is created through a patented, water-based process that combines vitamin C and calcium. Ester-C® is non-acidic and should be well-tolerated by those who are sensitive to the acidic nature of ascorbic acid.

FUNCTIONS

Vitamin C: Vitamin C (ascorbic acid) has numerous biological functions. It is essential for the synthesis of collagen and glycosaminoglycans, which are the building materials of all connective tissues, such as skin, blood vessels, tendons, joint cartilage and bone. Vitamin C is the required coenzyme for two groups of enzymes that catalyze the crosslinking of collagen fibers - lysyl hydroxylases and prolyl hydroxylases. As such, vitamin C is essential for normal wound healing and capillary health.

Vitamin C participates in the biosynthesis of carnitine, serotonin, and certain neurotransmitters, including norepinephrine.

Vitamin C is among the most powerful antioxidants in humans and animals. It is a water-soluble, chainbreaking antioxidant that reacts directly with superoxide, hydroxyl radicals, and singlet oxygen. Laboratory studies show that vitamin C completely protects lipids in plasma and low-density lipoprotein (LDL) against atherogenic peroxidative damage. In addition, vitamin C interacts with glutathione and alpha-lipoic acid, and regenerates vitamin E.

The antioxidant functions of vitamin C appear to have clinical significance in providing protection from free radical damage to the eyes, lungs, blood and the immune system.

Vitamin C is absorbed in the small intestine by a sodium-dependent transport process that is intake dependent. At normal dietary intakes of 60 to 100 mg, up to 80 or 90% of the vitamin C is absorbed. At higher intakes, absorption becomes less efficient. Absorption efficiency and vitamin C utilization may be greatly enhanced during conditions of physiological stress, such as trauma or infection. Maximal absorption is attained by the ingestion of several doses spaced throughout the day rather than in one, larger dose.

Citrus Bioflavonoids: The white layer of citrus peels is especially rich in the bitter tasting flavanones hesperidin and naringin. Citrus juices, however, are low in bioflavonoids. Like most flavonoids, those from citrus are free radical scavenging antioxidants that work in synergy with vitamin C. Ester-C® Complex provides 100 mg of a concentrated bioflavonoid complex.

INDICATIONS

Ester-C® Complex may be a useful dietary supplement for those who wish to supplement their diet with this complementary blend of a unique, non-acidic form of vitamin C and citrus bioflavonoids.

FORMULA (EST)

Each Vegetarian capsule contains:	
Vitamin C	490mg
(as Ester-C®** calcium ascorbate)	-
Calcium	60mg
(as Ester-C®** calcium ascorbate)	-
Citrus Bioflavonoid complex	100mg
(includes hesperidin and naringen)	-

**Ester-C® is a licensed trademark of INTER-CVAL Corporation. Manufactured under U.S. Patent No. 4,822,816.

1

Ester-C® Complex

Vitamin C as Calcium Ascorbate with Citrus Bioflavonoids

SUGGESTED USE

Adults take 1 to 2 capsules daily or as directed by a healthcare professional.

SIDE EFFECTS

No adverse side effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Ameer B, Weintraub RA, Johnson JV, et al. Flavanone

absorption after naringin, hesperidin, and citrus administration. Clin Pharmacol Ther 1996 Jul;60(1):34-40. Ballmer PE, Reinhart WH, Jordan P, Bühler E, Moser UK, Gey KF. Depletion of plasma vitamin C but not of vitamin E in response to cardiac operations. J Thorac Cardiovasc Surg 1994:108:311-320.

Barabás J, Nagy E, Degrell I. Ascorbic acid in cerebrospinal fluid--A possible protection against free radicals in the brain. Arch Gerontol Geriatr 1995;21:43-48.

Bendich A, Langseth L. The health effects of vitamin C supplementation: A review. J Am Coll Nutr 1995;14:124-136.

Berkarda B, Koyuncu H, Soybir G, Baykut F. Inhibitory effect of

Hesperidin on tumour initiation and promotion in mouse skin. Res Exp Med (Berl) 1998 Aug;198(2):93-9. Beyer RE. The role of ascorbate in antioxidant protection of biomembranes: Interaction with vitamin E and coenzyme Q. J Bioenerg Biomembr 1994;26:349-358.

Bielory L, Gandhi R. Asthma and vitamin C. Ann Allergy 1994;73:89-96.

Bush MJ, Verlangieri AJ. An acute study on the relative gastro-intestinal absorption of a novel form of calcium ascorbate. Res Commun Chem Pathol Pharmacol 1987;57:137-140.

Cathcart RF, III. Vitamin C in the treatment of acquired immune deficiency syndrome (AIDS). Med Hypotheses 1984;14:423-433.

Cunningham JJ, Mearkle PL, Brown RG. Vitamin C: An aldose reductase inhibitor that normalizes erythrocyte sorbitol in insulin-dependent diabetes mellitus. J Am Coll Nutr 1994;13:344-350.

Edwards DJ, Bernier SM. Naringin and naringenin are not the primary CYP3A inhibitors in grapefruit juice. Life Sci 1996;59(13):1025-30.

Eriksson J, Kohvakka A. Magnesium and ascorbic acid supplementation in diabetes mellitus. Ann Nutr Metab 1995;39:217-223.

Faruque MO, Khan MR, Rahman M, Ahmed F. Relationship between smoking and antioxidant nutrient status. Br J Nutr 1995;73:625-632.

Franke AA, Cooney RV, Custer LJ, et al. Inhibition of neoplastic transformation and bioavailability of dietary flavonoid agents. Adv Exp Med Biol 1998;439:237-48.

Frei B. Reactive oxygen species and antioxidant vitamins: Mechanisms of action. Am J Med 1994;97 Suppl. 3A:5S-13S.

Galati EM, Trovato A, Kirjavainen S, Forestieri AM. Biological effects of hesperidin, a Citrus flavonoid. (Note III): antihypertensive and diuretic activity in rat. Farmaco 1996 Mar;51(3):219-21.

Gale CR, Martyn CN, Winter PD, Cooper C. Vitamin C and risk of death from stroke and coronary heart disease in cohort of elderly people. BMJ 1995;310:1563-1566.

Gerster H. Antioxidant vitamins in cataract prevention. Z Ernahrungswiss 1989;28:56-75.

Ghosh SK, Ekpo EB, Shah IU, Girling AJ, Jenkins C, Sinclair AJ. A double-blind, placebo-controlled parallel trial of vitamin C treatment in elderly patients with hypertension. Gerontology 1994;40:268-272.

Green MHL, Lowe JE, Waugh APW, Aldridge KE, Cole J, Arlett CF. Effect of diet and vitamin C on DNA strand breakage in freshly-isolated human white blood cells. Mutat Res DNAging Genet Instability Aging 1994;316:91-

Ester-C® Complex

Vitamin C as Calcium Ascorbate with Citrus Bioflavonoids

102.

Guilhou JJ, Fevrier F, Debure C, et al. Benefit of a 2-month treatment with a micronized, purified flavonoidic fraction on venous ulcer healing. A randomized, double-blind, controlled versus placebo trial. Int J Microcirc Clin Exp 1997;17 Suppl 1:21-6.

Hennekens CH. Antioxidant vitamins and cancer. Am J Med 1994;97 Suppl. 3A:2S-4S.

Jariwalla RJ, Harakeh S. Ascorbic Acid and AIDS: Strategic Functions and Therapeutic Possibilities. In: WATSON RR, ed. Nutrition and AIDS. Boca Raton: CRC Press, Inc. 1994:117-139.

Johnston CS, Yen M-F. Megadose of vitamin C delays insulin response to a glucose challenge in normoglycemic adults. Am J Clin Nutr 1994;60:735-738.

Kawaguchi K, Mizuno T, Aida K, Uchino K. Hesperidin as an inhibitor of lipases from porcine pancreas and Pseudomonas. Biosci Biotechnol Biochem 1997 Jan;61(1):102-4.

Khaw K-T, Woodhouse P. Interrelation of vitamin C, infection, haemostatic factors, and cardiovascular disease. BMJ 1995;310:1559-1563.

Kim DH, Jung EA, Sohng IS, et al. Intestinal bacterial metabolism of flavonoids and its relation to some biological activities. Arch Pharm Res 1998 Feb;21(1):17-23

Kodama M, Kodama T. Vitamin C and the genesis of autoimmune disease and allergy. In Vivo 1995;9:231-238.

Kodama M, Kodama T, Murakami M. Autoimmune disease and allergy are controlled by vitamin C treatment. In Vivo 1994;8:251-258.

Lambev I, Belcheva A, Zhelyazkov D. Flavonoids with antioxidant action (naringin and rutin) and the release of mastocytic and nonmastocytic histamine. Acta Physiol Pharmacol Bulg 1980;6(2):70-5.

Lehr H-A, Frei B, Arfors K-E. Vitamin C prevents cigarette smoke-induced leukocyte aggregation and adhesion to endothelium in vivo. Proc Natl Acad Sci USA 1994;91:7688-7692.

Miyake Y, Yamamoto K, Tsujihara N, Osawa T. Protective effects of lemon flavonoids on oxidative stress in diabetic rats. Lipids 1998 Jul;33(7):689-95.

Mukhopadhyay CK, Chatterjee IB. Free metal ion-independent oxidative damage of collagen. Protection by ascorbic acid. J Biol Chem 1994;269:30200-30205.

Paolisso G, Balbi V, Volpe C, et al. Metabolic benefits deriving from chronic vitamin C supplementation in aged non-insulin dependent diabetics. J Am Coll Nutr 1995;14:387-392.

Sauberlich HE. Pharmacology of vitamin C. Annu Rev Nutr 1994;14:371-391.

Seddon JM, Ajani UA, Sperduto RD, et al. Dietary carotenoids, vitamins A, C, and E, and advanced agerelated macular degeneration. Eye Disease Case-Control Study Group. JAMA 1994;272:1413-1420.

Tanaka T, Makita H, Kawabata K, et al. Chemoprevention of azoxymethane-induced rat colon carcinogenesis by the naturally occurring flavonoids, diosmin and hesperidin. Carcinogenesis 1997 May;18(5):957-65. Verlangieri AJ, Fay MJ, Bannon AW. Comparison of the anti-scorbutic activity of L-ascorbic acid and Ester C in the non-ascorbate synthesizing Osteogenic Disorder Shionogi (ODS) rat. Life Sci 1991;48(23):2275-81. Winkler BS, Orselli SM, Rex TS. The redox couple between glutathione and ascorbic acid: A chemical and physiological perspective. Free Radic Biol Med 1994;17:333-349.

For more information on Ester-C® Complex visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by Douglas Laboratories 600 Boyce Road Pittsburgh, PA 15205 800-245-4440 douglaslabs.com



You trust Douglas Laboratories. Your patients trust you.